

## Microfacies and depositional basin analysis of Cretaceous deposits at Korkouh area

Narjes Naraki<sup>1,\*</sup>, Mohammad Dastanpur<sup>2</sup>, Mohammad-Javad Hasani<sup>2</sup>

1- Graduate student of Stratigraphy and Paleontology, Institute of Higher Education of Kerman, Iran.

2- PhD of Stratigraphy and Paleontology, Institute of Higher Education of Kerman, Iran.

\* Corresponding Author: Naraki.narjes@gmail.com

### Abstract

In order to microfacies studies of the Cretaceous deposits, a section called Korkouh was selected in Kerman province in Central Iran area. The Korkouh is on the Bidou Formation with angular unconformity and underlain by paleogene conglomerates with an erosional surface. The studied section basically contains lime and dolomitic limestone with the thickness of 146 meters with new fossils such as foraminifera, bivalves and bryozoans. In general, foraminifera obtained from Korkouh section indicate the existence of sediments in the age of Albian and Cenomanian. Based on microfacies studies on the section three facies belts have been identified in Korkouh section that include intertidal, lagoon and inner ramp. Studying of identified facies and comparing them with the present sedimentary environment showed that Korkouh outcrop facies have been deposited.

**Keywords:** Cretaceous; Korkouh Section; Microfacies; Biostratigraphy; Foraminifera; Kerman, Intertidal, Lagoon.

### 1- Introduction

Recognition of facies is very important in stratigraphy layers. Facieses are studied in both microscopic and in macroscopic scales. In this study, microfacies are named based on the microfacies of Flügel's standard (2004), and according to available lithological and paleontological features, we tried each of microfacieses compare the facies belts of Flügel's model as much as possible. In general, lithological and paleontological characteristics of a sedimentary facies have to be specified, so that facies are easily separated from each other. Differences in environmental factors (sedimentology and biological), are the main reasons for the differences in microscopic facies, which in detail including the impact of organisms, water energy, oxygen level, water depth, light intensity, salinity, water temperature, arrival rate of clastic material and substrate type (Flügel, 2004). The studied cross-

section in Korkouh is 35 Km South East of Kerman (Fig. 1).

### 2- Materials and Methods

In order to study microbiostratigraphy and depositional environments of the Cretaceous deposits in the South East of Kerman, at first geological map of the study area were identified for selecting the section, then site visits and sampling of this area was carried out. After preparation of the samples were studied. Systematic sampling and sampling intervals at selected section, was used on average, between one and three meters.

A total of 20 samples at harvest field was collected which prepared thin layers of them. Then, in the laboratory were studied by polarizing microscope and then identify foraminifera available, microfacies of each sample was determined. The microfacies named using Dunham (1962). Sedimentary facies diagnosis and determine the environment in

accordance with the model proposed by Flügel (2004) has been made. According to field observations, accurate sampling, laboratory studies, data analysis and final conclusion, the article was written.




Figure 1) Location map of the study area.

### 3- Geological setting

The Korkough section is located in the 35 kilometers SE of Kerman in longitude  $57^{\circ} 25'$  and latitude  $30^{\circ} 05'$ . Geologically, the province with about  $183,714 \text{ Km}^2$  area situated in the southeastern region of Iran and in the provinces of Hormozgan, Sistan and Baluchestan, Fars, Yazd and Southern Khorasan which encompasses nearly 11% of the total area of the country.

#### 2.1- Stratigraphy

The Cretaceous limestone and dolomitic limestone layers have angular unconformity on Willow formation and is covered by an erosional surface

and it is in turn is covered by Paleogene conglomerates. The thickness of the harvest section is 146 m. The study of layers aligned along the NW–SE direction. The total of 20 representative samples were collected for biostratigraphy of microfacies studies (Fig. 2).




Figure 2) Stratigraphic column studied region.

#### 2.2- Lithology

The limestone and dolomitic limestone in the study area are comprised of following layers:

- The 116.5 m gray limestone containing fossils of Bryozoa, in the bivalve, planktonic and benthic foraminifera and the detrital components (Intraclasts).
- The 29.5 m dolomitic limestone containing fossils of Bryozoa, foraminifera and Intraclasts.

According to microfacies studies, facies have been identified in the Korkouh section. These microfacies are related to intertidal, lagoon and mid ramp.

#### **Intrataidal facies belt (A):**

The 17 samples are collected from this facies belt. One of the characteristics of this facies belt is the existence of dolomitized lime mudstone microfacies that is attributed to intertidal with no allochemical components which is composed of lime mud that locally show bird's eye orthochem. The Secondary dolomitization is one of the diagenetic phenomena happening in such microfacies (Fig. 3).




Figure 3) Microphotograph of intertidal dolomitized lime-mudstone facies.

#### **Lagoon (B):**

This facies belt includes the following Microfacies:

Bioclast Intraclast wackestone / packstone, bioclast intraclast wackestone, Dolomitized bioclast packstone, bioclast wackestone/packstone, Bioclast packstone, bioclast wackestone, Dolomitized bioclast Intraclast packstone/wackestone, with thickness of 129.5 m.

#### **Bioclast packstone microfacies (B1):**

This microfacies contains 40-50 % of bioclasts including *Charentia* sp., *Cyclammina* sp., *Praechrysalidina* infracretacea, *Nezzazata* sp., *Minouxia lobata* is attributed to lagoon zone. It is worth noting that such diagenetic phenomena calcite veins in the desired microfacies (Fig. 4).




Figure 4) Microphotograph of bioclast packstone facies belt lagoon.




Figure 5) Microphotograph of bioclast wackestone facies belt lagoon.

#### **Bioclast wackestone microfacies (B2):**

According to main allochemicals 40% of *Nummofallotia* apull, *Nezzazata* sp., *minouxia*

lobata, Textularia sp., Hemicyclamminasp., Rzehakina sp., Praechrysalidina infracretacea, Pseudolituonella sp., Cuneolinapavonia, Quinqueloculina sp. with traces of bryozoas, echinoderms pieces, ostracoda. Regarding to the mentioned cases, the following microfacies are in the central part with less energy of lagoon environment (Fig. 5).

#### **Bioclast Intraclast wackestone microfacies (B3):**

Skeletal components of this microfacies contain 15% benthic foraminiferas such as Nezzazata sp., Pseudocyclammina lituus, Nummofallotia sp., Textularia sp., Praechrysalidina infracretacea with bivalve debris 4%, and non-skeletal components contain 20% intraclast pieces that are not well rounded and sorted due to low energy of environment (Mousavi-Harami, 2000; Fig. 6).




Figure 6) Microphotograph of bioclast intraclast wackestone facies belt lagoon.

#### **Bioclast Intraclast wackestone/ packstone microfacies (B4):**

The main components of this microfacies are Minouxia lobata, Textularia sp., Praechrysalidina infracretacea, Nezzazata sp., Charentia cuvillieri, gastropod and Bryozoa, intraclast pieces in calcareous ooze orthochem. Porcelaneous walled foraminifera presence of

water is sign of peaceful environment and with limited rotation (Hottinger, 1997; Racey, 2001) (Fig. 7).




Figure 7) Microphotograph of bioclast intraclast wackestone/ packstone facies belt lagoon.

#### **Bioclast wackestone/packstone microfacies (B5):**

This microfacies contains of Nezzazata sp., Cuneolina pavonia, Nummofallotia sp., Charentia sp., Textularia sp., Praechrysalidina infracretacea, Minouxia lobata, Miliolid and of gastropod and Bryozoa in lime orthochem (Fig. 8).

#### **Dolomitized bioclast packstone (B6):**

Its main components are Quinqueloculina sp., Rzehakina sp., Miliolid, Cuneolina pavonia, textularia sp., Nummofallotia sp., and Bryozoa in micritic orthochem. Rocks containing Miliolid are attributed to lagoon zone (Geel, 2000). The secondary dolomitization is one of the diagenetic phenomena that occurred in such microfacies (Fig. 9).

#### **Bioclast Intraclast packstone microfacies (B7):**

The main components of this microfacies are Nezzazata sp., Rzehakina sp., Minouxia lobata, Quinqueloculina sp., Pseudocyclammina sp.,

Bryozoa, and intraclast in lime ortochem (Fig. 10).




Figure 8) Microphotograph of bioclast wackestone/packstone facies belt lagoon.




Figure 9) Microphotograph of dolomitized bioclast packstone facies belt lagoon.

#### **Dolomitized bioclast packstone/ wackestone (B8):**

This microfacies is composed of *Quinqueloculina* sp., Miliolid, *Cuneolinapavonia* and Bryozoa in lime ortochem. Rocks containing Miliolid are attributed to lagoon zone (Geel 2000; Fig. 11).




Figure 10) Microphotograph of bioclast intraclast packstone facies belt lagoon.




Figure 11) Microphotograph of dolomitized bioclast packstone/wackestone facies belt lagoon.


#### **4- Discussion**

The main allochem in microfacies of the lagoon is small benthic order of Miliolina (Miliolides). This is located in the inner ramp of microfacies lagoon. The facies with benthic shells along the lagoon of vans, the producer of the low energy environment, upper photic zone and sat down the bottom of the shallow lagoon commeg inner ramp section. (Romero et al., 2002) with more close to the barrier and the increase of energy

and environment, the value of the wackestone facies of micritic costs and value added sparite packstone facies. Presence of porcelaneous foraminifera implies high salinity characteristics of the environment (Flügel, 2004). Furthermore, the abundance of Miliolid may be attributed to the lagoon (Penney and Racey, 2004).

#### **Mid ramp facies belt (C):**

This facies has thickness of 13.5 m. The existence of bioclast wackestone microfacies is one of the characteristics of this facies.



Microfacies in such facies includes benthic foraminifera associated with Planktonic foraminifera which may implies its formation in restricted lagoons in pools to the open sea. It is confirmed by presence of the fauna of both open sea and pool.

#### **Bioclast Wackestone microfacies (C1):**

This microfacies contains Benthic foraminifera including *Charentia* sp., *Textularia* sp., with Planktonic foraminifera such as *Globotruncana* cf. *helvetica*, *Globotruncana* sp. and *Ostracoda* (Fig. 12).




Figure 12) Microphotograph of bioclast wackestone facies belt Mid ramp.

#### **Depositional model:**

Since the sequence of carbonate facies result of changes in the environment through time (Tucker and Wright, 1990), it is possible to identify microfacies and their formation conditions and interpret their depositional environment and.

The described facies belts of tidal, lagoon and the open sea for microfacies have been emplaced in the shallow carbonate platform. The observed vertical changes in microfacies represents a lack of re-deposited sediments and symptoms caused by a sudden change, for example turbidites represent the slope of the

carbonate platform (Burchette and Wright, 1992). The facieses of carbonate sediments in the reef barrier formed a carbonate ramp, because expansion is limited to reef ramp feature (Einsele, 2000).

According to environmental studies on the proposed model can be set up in each of the horizons and facies belts in all positions during section conclude that early cutting (horizons 1 to 7) to constantly advance and recession of the sea(s). Half of the section to the next (horizons 8 to 16), and at the end of a cycle, steadily cutting back the advance and recession encounter sea (Fig. 13).




Figure 13) Facies belts distribution and facies models in studied region.

**4- Conclusion**

- 1- The 146m Korkouh outcrop (SE Kerman) lithology is mainly limestone and dolomitic. In this study 21 genus and species of foraminifera on the Korkouh outcrop have been identified.
- 2- According to microfacies studies, facies have been identified in Korkouh section and Theseer microfacies are related to

intertidal, lagoon and mid ramp in the carbonate platform.

- 3- The existence of fossil foraminifera genus and species accumulation index and age region of the Albian-Cenomanian show the emergence and spread of *Minoxia lobata*, *Nezzazata conica*, *Cuneolina pavonia*, *Pseudocyclamina* sp., could be confirmed this age of the study area.

- 4- Korkouh level with angular unconformity is on the Bidou formation and covered with an erosional surface by paleogen econgglomerates.

## References:

- Aryaie, A. A., Glumaky Rabati, H. 2007. Microfacies of Upper Cretaceous deposits in the south of Valley Golbaf Home Khatun. 11<sup>th</sup> Conference of Geological Society of Iran. Ferdowsi University of Mashhad, Iran.
- Al-Ali, S. M., Phaiz-Niya, S., Esmail, A., Shahrooi, A. 2007. Sarvak Microfacies and depositional environment in the North West Behbahan (anticline Bangestan), 11<sup>th</sup> Conference of Geological Society of Iran, Ferdowsi University of Mashhad, Iran.
- Asilian, H., Lasemi, Y. 2007. Ilam Formation sedimentary environments and sequence stratigraphy in the North West Gulf, South West of Iran, 11<sup>th</sup> Conference of Geological Society of Iran, Ferdowsi University of Mashhad, Iran.
- Omidvar, M., Sajadi, F. 2011. Biostratigraphy and determine the Sarvak and the lower part of the Ilam Formation in oil fields and water maroon Teymur. 15<sup>th</sup> Congress of Geological Society of Iran, Tarbiat Moallem University, Tehran, Iran.
- Iran-Nejad, M., Ghotby, M. R., 2010. Microbiostratigraphy Cretaceous deposits in North Raver (mountain section markesh). Regional Conference on Geology plateau of Iran, Islamic Azad University of Kerman, Iran.
- Bord-Bart, A., Gorgij, M. N., Mousavi-Harami, R., Najafi, M. 2010. Microfacies and depositional environment of the Upper Cretaceous deposits in the North West Nehbandan. The 29<sup>th</sup> meeting of Earth Sciences, Geological Survey of Iran.
- Rahiminejad, A. H., Vaziri-Moghadam, H., Syrfyan, A., 2006. Biostratigraphy and Microfacies Sarvak Traditional wells, Research journal Science: 23, 87–104 Isfahan.
- Satari, P., Mousavi, M. R., Shah-Bakhsh Majbour, A. 2012. Facies, sedimentary environments and sequence stratigraphy Sarvak the anticline cumin (Gulf Coast). The 6<sup>th</sup> National Conference of Geology, Payam-e-Noor University, Kerman, Iran.
- Kazem-Zadeh, M. H., Ghasemi-Nezhad, A., Vaziri-Moghadam, H., 2011. Biostratigraphy and depositional environment Sarvak study in Ahvaz oil field. The 15<sup>th</sup> Congress of Geological Society of Iran, Tarbiat Moallem University, Tehran, Iran.
- Kavousi, M. A. 2005. Facies and depositional environments of the Upper Cretaceous deposits north East of Kalar Dasht. 15<sup>th</sup> Congress of Geological Society of Iran, Tarbiat Moallem University, Tehran, Iran.
- Karimi Bavand-Pour A. R. 1994. Microfacies and depositional environment Sarvak in Khuzestan (Aghajari well). The 8<sup>th</sup> Congress of Geological Society of Iran, Shahrood.
- Mahani-Pour, A. 1383. Biostratigraphy and paleoecology study Upper Cretaceous deposits in the Mountain West Badamouyeh Kerman. MSc thesis, University of Kerman, 281 p.
- Maleki Khimeh-Sari, S., Khosrotehrani, Kh., Asilian-Mahabadi, H., Lasemi, Y. 2009. Formation outcrop of sedimentary environments and sequence stratigraphy, sectioning surface and section Consisted of Bidkhon and Khartang two and started ground fire west. Journal of Applied Geology: 4, 298–287.
- Nozari-Pour, B., Vaziri, S. H., Taherpour Khalilabad, M., Sedaghat, M. A. 2011. Biostratigraphy Kazhdumi Formation in the stratigraphic section Scholl North West of Shiraz. 2<sup>nd</sup> Congress of Geological Society of Iran, University of Mashhad, Iran.
- Burchette, T. P., Wright, V. P. 1992. Carbonate ramp depositional systems. Sedimentary Geology: 79, 3–35.
- Dunham, R. J. 1962. Classification of Carbonate Rocks According to Depositional Texture. In, W.E. Hamm (Ed.), Classification of Carbonate Rocks, A Symposium. American Association of Petroleum Geologists: 108-121.
- Einsele, G. 2000. Sedimentary Basin: Evolution, Facies and sediment Budget. 2nd ed., Springer-Verlag: 292p.
- Flügel, E. 2004. Microfacies Analysis of Carbonate Rocks: Analysis, Interpretation and Application. Springer-Verlag: 976p.

Geel, T. 2000: Recognition of stratigraphic sequences in carbonate platform and slope deposits: empirical models based on microfacies analysis of Paleogene deposits in southeastern Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology*: 155, 211–238.

Romero, J., Caus E., Rossel, J. 2002. A model for the palaeoenvironmental distribution of larger foraminifera based on Late Middle Eocene deposits on the margin of the south Pyrenean basin (SE Spain): *Palaeogeography, Palaeoclimatology, Palaeoecology*. 179: 43–56.

Tucker, M. E., Wright, V. P. 1990. *Carbonate sedimentology*. Blackwell, Oxford. 482 p.

**Received: 13 September 2015 / Accepted: 01 December 2015 / Published online: 05 December 2015**

#### EDITOR-IN-CHIEF:

##### *Dr. Vahid Ahadnejad:*

Payame Noor University, Department of Geology, PO BOX 13395–3697, Tehran, Iran.  
E-Mail: edchief@jtethys.org

#### EDITORIAL BOARD:

##### *Dr. Jessica Kind:*

ETH Zürich Institut für Geophysik, NO H11.3, Sonneggstrasse 5, 8092 Zürich, Switzerland  
E-Mail: jessica.kind@erdw.ethz.ch

##### *Prof. David Lentz*

University of New Brunswick, Department of Earth Sciences, Box 4400, 2 Bailey Drive  
Fredericton, NB E3B 5A3, Canada  
E-Mail: dlentz@unb.ca

##### *Dr. Anita Parbhakar-Fox*

School of Earth Sciences, University of Tasmania, Private Bag 126, Hobart 7001, Australia  
E-Mail: anitap1@utas.edu.au

##### *Prof. Roberto Barbieri*

Dipartimento di Scienze della Terra e Geoambientali, Università di Bologna, Via Zamboni 67 – 40126, Bologna, Italy

E-Mail: roberto.barbieri@unibo.it

##### *Dr. Anne-Sophie Bouvier*

Faculty of Geosciences and Environment, Institut des sciences de la Terre, Université de Lausanne, Office: 4145.4, CH–1015 Lausanne, Switzerland  
E-Mail: Anne-Sophie.Bouvier@unil.ch

##### *Dr. Matthieu Angeli*

The Faculty of Mathematics and Natural Sciences, Department of Geosciences, University of Oslo  
Postboks 1047 Blindern, 0316 OSLO, Norway  
E-Mail: matthieu.angeli@geo.uio.no

##### *Dr. Miloš Gregor*

Geological Institute of Dionys Stur, Mlynska Dolina, Podjavorinskej 597/15  
Dubnica nad Vahom, 01841, Slovak Republic  
E-Mail: milos.gregor@hydrooffice.org

##### *Dr. Alexander K. Stewart*

Department of Geology, St. Lawrence University, Canton, NY, USA  
E-mail: astewart@stlawu.edu

##### *Dr. Cristina C. Bicalho*

Environmental Geochemistry, Universidade Federal Fluminense – UFF, Niterói–RJ, Brazil  
E-mail: ccbicalho@gmail.com

##### *Dr. Lenka Findoráková*

Institute of Geotechnics, Slovak Academy of Sciences, Watsonova 45, 043 53 Košice, Slovak Republic  
E-Mail: findorakova@saske.sk

##### *Dr. Mohamed Omran M. Khalifa*

Geology Department, Faculty of Science, South Valley University, Qena, 83523, Egypt  
E-Mail: mokhalifa@svu.edu.eg

##### *Prof. A. K. Sinha*

D.Sc. (Moscow), FGS (London). B 602, Vigyan Vihar, Sector 56, GURGAON 122011, NCR DELHI, Haryana, India  
E-Mail: anshuksinha@gmail.com