

## Recognition of Reservoir Continuity and Reservoir Compartmentalization Using Oil Finger Printing In One of Iranian Oil Region in the Persian Gulf

Idin Shahrezaei<sup>1\*</sup>, Mohammad Reza Kamali<sup>2</sup>, Amirreza Abadichalaksaraee<sup>1</sup>

1- Department of Geology, Faculty of Sciences, Islamic Azad University, Science and Research Branch, Tehran, Iran.

2- Faculty of Research and Development in Upstream Petroleum Industry-RIPI.

\* Corresponding Author: idn\_sha@yahoo.com

### Abstract

Salman and Resalat oilfields are such marine oilfields that are in vicinity of each other. There is Resalat oilfield in distance about 80-90 kilometers from Lavan Island in Persian Gulf. This oilfield is located in 25 kilometer southwest in Reshadat oilfield. Salman oilfield is located in vicinity of water border line of United Arab Emirate in 140 kilometers of south of Lavan Island. This oilfield is remarkable because of its reservoir value and also it is common between Iran and United Arab Emirate. Surmeh reservoir structure is an oily reservoir that it's common between two oilfields. In order to study of geochemistry features, oil samples of Salman and Resalat's oilfield analyzed by using usual geochemistry techniques such SARA test, gaseous chromatography techniques (GC) and gaseous chromatography – mass spectrograph (GC-MS). According to results, including of distribution of values in formed hydrocarbon cuttings of these oils, study of Biomarkers in Terpens and tabloid steranes group from saturation cutting, we can say that hydrocarbons in Surmeh reservoir in Resalat and Salman oilfield show Paraffinic oil features and kerogen type II that all of these indicate organic material with marine origin. The geochemical study of parent rock in these oilfields shows that the active oil system comes from late Cretaceous and the reservoirs were fed from common hydrocarbon system. The gathering hydrocarbons at mentioned oilfields, also, were produced by carbonate source rock with a little shale in revival conditions. The parameters of mature biomarkers in these oilfields show that in border of two oil-forming windows it gathered the suitable organic material for producing good oil. Thus, according to analyses, there is integration between Surmeh reservoir in Resalat and Salman oilfield and their oil come from one resource.

**Keywords:** Reservoir Geochemistry, Surmeh structure, Integration, Resalat oilfield, Salman oilfield, Biomarker.

### 1- Introduction

Determination and evaluation of one area in terms of kind of oil in reservoir, the effect of secondary processes and determination of their probably source stones and vertical and horizontal continuity of oil reservoir together, all of them, are important for exploration targets of experts from the past. Persian Gulf zone is one of the strategic and important zones in Iran and there are more than 3 billion barrels of

recoverable oil in this zone. Geochemistry evaluation of the oil of this area can be important from investigation, utilization and development point of view (Fig.1, Rabbani, 2014).

Therefore, with the study of probably oil resources in available reservoir we can act with more sure in search and use of hydrocarbon resources and in this regard we can planning for oilfields that have not been discovered by now (Stratigraphic oil trap). The studying reservoir

stones in Resalat and Salman oilfield are in Surmeh reservoir structure which they have mainly carbonate lithology (Rabbani, 2013). In this regard, recognition of the oil in producing layers which are in side of them horizontally, study of biological resources and signs by oil

fingerprint technique, evaluation of production or reservoir continuity and secondary recovery by oil fingerprint effect, recognition of local and regional probably obstacles of vertically and horizontally of hydrocarbons in reservoir, all of them are next station in this process.




Figure1) Location of Salman and Resalat Oilfield in Persian Gulf.

**2- Research methodology**

To determine of Biomarkers and kinds of them (Bordenave, 1993) and find out information about distribution of normal Alkanes and Isoprenoids, first, by use of samples of oils and resource stones, have been done experiments such as GC and GC-MS on the samples. Secondly, by using of oil fingerprint find the information about index biological signs of sedimentary environment, the mature degree of organic material and oil, analysis option like leaching or bacterial breakdown, conformity of oil to oil and oil to resource rock (Waples, 1991), and finally by using these information we can achieve our goal and generally manage and use reservoirs optimize. Also, by use of FIS new analysis method or liquid inclusion stratigraphy which do it on samples of teeny of digging and cores, we can draw the history of filling the reservoir and the plan of stratigraphy of fluid chemical. Finally, we can answer many of production and investigations questions.

**3- Discussion**

To study and diagnosis of differences and similarities of hydrocarbonates in Surmeh reservoir, it has been used columnar chromatography techniques (SARA experiment). The studying oil samples by this method separate from each other. It has been determined different percent of hydrocarbon cuts. Hydrocarbon cuts are included of saturation hydrocarbons, aromatics, resins and asphaltene which this collection called pelar. Thus, the oils which have high saturation hydrocarbons have a very good quality. Increasing Aromatics make Aromatic oils. Heavily oils have more asphaltene and resin which in this kind of oil the quality is low.

Table1)seperated hydrocarbon cuts from Y1 and X5 samples.

Field/Reservoir	Saturate %	Aromatic %	Resin %	Asphaltene %
Resalat Y1	55	32	11	2
Salman X5	62.3	27.5	8.6	1.6

With drawing both oilfields on one chart, we compare two oilfields. As you see on Table 1 the quality of Salman oilfield is better than Resalat in Surmeh layer because oil of Salman

oilfield has better saturation hydrocarbons, it's aromatics and non-saturated hydrocarbons are low (Fig.2).




Figure 2) distribution of SARA cuts for X5 and Y1 samples.

Considering to placement of samples in triangle diagram (Fig.3), Salman oilfields samples are in Paraffin and Paraffin-Naphthenic but Resalat oilfield sample is in Paraffinic area (TissotandWelte, 1984). So by using the tree plot chart can find that the oil of these oilfields because of be paraffinic with high degree maturity is good in quality.

Table2) separated hydrocarbon cuts from studying oil samples.

Sample	Saturate (%)	Aromatics (%)	Resin (%)	Asphaltene (%)
Salman X1	45.8	29.6	21.1	3.5
Salman X2	35.1	40.4	21.5	2.9
Salman X3	40	34.7	21.8	3.5
Salman X4	34.8	36.2	26.2	2.8
Salman X5	62.3	27.5	8.6	1.6
Resalat Y1	55	32	11	2




Figure3) Triangle graph of formation cuts percent in testing samples in Surmeh layers in Salman and Resalat oilfield.

By use of ratio of odd Alkans to even Alkans with the Ph/nC17, Pr/nC18 and Pr/Ph ratios in star shape graph, also comparison of dispersion distribution of these ratios which are gain from analysis of gas chromatography on oil samples of any structure and oilfield, we can recognize sameness or different of hydrocarbones or propebly source ston fitures (Kaufman and Ahmad, 1990).

According to data (Table3) and drawing all of parameters on a star shape diagram (Fig. 4), we see that same distribution and suitable overlap to odd and even Alkans and non-cyclic isopronoeds show that these reservoirs are fed from same resource.

Table3) Parameters and ratios related to impregnating compounds from gas chromatographic analysis on oil samples in studying oilfields.

Well	Resalat	Salman X6	Salman X7
Formation	Arab C (Surmeh)	Arab D (Surmeh)	Arab A (Surmeh)
Pr/ph	0.59	0.54	0.59
Pr/nC17	0.18	0.19	0.2
Ph/nC18	0.34	0.37	0.36
C19/C20	1.06	1.04	1.04
C21/C22	1.12	1.12	1.12
C23/C24	1.06	1.06	1.04
C25/C26	1.08	1.08	1.08
C27/C28	1.13	1.12	1.12
C29/C30	1.12	1.11	1.05
C31/C32	1.17	1.14	1.14

To diagnosis of type and mature degree in organic matter, to determine of the environment of source stone in oilfield and comparison of oils together and to source stone, could use from different parameters of biomarkers. Biomarkers could be suitable patterns in evaluation of old environments (De Leeuw et al., 1995). Biomarkers use for lithology determination, special sedimentary age and environment source stone, source stone thermal maturity during producing process and also rational amount of producing oil to organic matter of producing gas in source Kerogen. These data have key role in oilfield forecasts model.




Figure 4) normal alkanes star shape diagram and isonoeds of samples studying

With use of Pri/n-c<sub>17</sub>, Phy/n-c<sub>18</sub> parameters, it could evaluate and determine the features of organic materials in this structures from the point of views of kind of Kerogen, process of maturity, probably process of alteration and the conditions of Sedimentation (Hunt, 1996).




Figure5)Ph/n C<sub>18</sub> diagram to Pr/n C<sub>17</sub> for the samples of Surmeh reservoir in Salman and Resalat oilfield.

In Resalat oilfield our samples show that the condition is suitable for oil. Maturity region and revive conditions show good quality and useful. Three samples of this oilfield are from Kerogen 2 which related to marine environments.

If Pr/Ph<1 shows that sedimentation of source stone happen in revive conditions (Didyk et al., 1978) (if Pr/Ph>1 thus the environment is oxygenic), thus the sedimentation environment is revival absolutely in Sarvak, Daryan and Surmeh Structure. In the meanwhile if Pr/Ph<0/8, the environment is a salty environment to hyper saline or is a carbonate environment and if Pr/Ph<0/3. By the way, more Fitan to Pristan confirm this which high

value of Fitan show revival environment and high value of Pristan show oxidic environment (Connan and Cassou, 1980). As we see the because of the samples placement in revival environment have  $Pr/Ph < 1$ .

Salman oilfield has exactly condition of Resalat oilfield. As you see, these samples in a special compress, when placement in Machur region, means that the samples have high maturity. When the oil is a high quality oil, it means that the condition was non oxygen condition. And finally the marin region tell us the kerogens were second type.




Figure 6)  $C_{35}/C_{34}$  Diagram in front of  $C_{29}/C_{30}$ .

The second method of determination of sedimental environmental and type of lithology is use of Hoopan and Homohoopan. The Homohoopan ratio  $C_{35}/C_{34}$  shows oxidation ratio of source stone environment. It suggest that as threshold of oxidic and nonoxidic environment, ratio of  $C_{35}$  and  $C_{34}$  homopans becomes one. If this ratio were less than one confirm that the sedimental environment is revival and if were more than one then we find out the sedimental environment is oxidic (Connan and Cassau, 1980).

As we see, our oil has consisted in many sedimental environment which include of carbonates and shales. By the way, as it placement in low of diagram, we can find that the condition was a revival and this situation is a marine position exactly. (Fig. 6).

In Resalat oilfield we study  $Y_1$  and  $Y_2$  samples, these samples come from this oilfield and Surmeh layer. This area shows a sedimentary

environment in kind of shale and shale-carbonate, because of a shale layer. The conditions of this oilfield is same as Salman oilfield which is a revival situation.

With use of  $T_s/(T_s+T_m)$  and  $Dia/(Dia+Reg)$  we can study the condition of sedimentary environment, by the way we can study if the environment acidic and alkaline pH or environment revival and oxidic EH or mature thermal also has effect on organic matter. We studied all of these subjects and found out the informations (Moldovan et al., 1994).

$T_s$  is a couple of hopans which there is in all samples of them.  $T_m$  is a biological type that because of diagenesis turn to  $T_s$  gradually. Accordingly  $T_s$  is a hopan thermo type. Diasterns which are arranged strans, are produced in catagenesis and diagenesis phase also.




Figure 7)  $T_s/(T_s + T_m)$  diagram in  $Dia / (dia+reg)$  sterane for all Surmeh reservoir samples.

The samples in Salman oilfield show revival environment in kind of shale sedimentary environment. This method is used for topographic diagnosis of course and is not accurate for lithology of resource stones. Also the samples were in thermo maturity too much which could be burn the samples and it has negative effect in oil quality.

Because of in marine environment both of Salman and Resalat oilfields are in vicinity of each other, the samples of Resalat oilfield has revival conditions and are affected thermal maturity (Fig. 7).

Sterane diagram in peristanfitan ratio is one of drawing methods of sedimentary environment.

According to evaluations,  $C_{27}$  Sterane is index of marine environments,  $C_{28}$  steranes is index of Lake Environment and  $C_{29}$  Steranes are indexes of continental environment (Peters et al., 2005). The resource of  $C_{29}$  steranes are usually from land plants which find in kerogens of type III, but the resource of  $C_{27}$  steranes are from marine fitoplanktones and  $C_{28}$  steranes are from lake algae and find in crogenes of type II and I respectively (Ekweozoret al.,1979).




Figure 8) Sterane  $C_{29}/C_{27}$  diagram in Pr/Ph for all samples of Surmeh reservoir of Salman and Resalat oilfield.

According to figure 7 in Salman oilfield we can see revival sedimentary environment in surmeh structure while there is in a algae region we find that oil is collecting in marine environment. So, Surmeh reservoir has placement in sea and oil marine crogenes are type of II. For samples the ratio of Peristan to fitian is less than one and the ratio of Sterane of  $C_{29}$  to  $C_{27}$  is between one to two that more Sterane of  $C_{27}$  show a marine environment.

The situation in Resalat oilfield is exactly same to Salman oilfield in Surmeh structure. When the oil of these two oilfield placement in revival and marine environment and of course to have algae and more esterane  $C_{27}$ , show that oil samples of two Salman and Resalat oilfield have same condition and quality. This figure could be one the signs that are from one resource. In this case, both oilfields are fed from one resource stone (Fig. 8).

The thermal mature shows the degrees of reactions from heat that sedimentary organic materials turns to oil. In this section, for determination of thermal maturity have been

used molecule parameters based on distribution and ratio of specific biological signs. Some of special ratios of saturation and aromatic biological signs are considered as most important thermal maturity (Peters et al., 2005). In general, organic material are divided to immaturity, maturity and over maturity, based on their relation to oil window. For finding the maturity of crudeoil, extracted bitomenes and their collection time the standard diagram of change rates  $C_{32} - \text{Hopane } 22S/(22S + 22R)$  in  $C_{29} - \text{sterane } 22S/(22S + 22R)$  can be used (Ourisson et al.,1984).

Mostly, the samples are between two oil-forming windows and have good quality.

There are two samples also in Resalat oilfield which are same condition to Salman oilfield samples in Surmeh layer. These features tell us the oil is high quality which gets a thermal maturity well. Finally, we can see thermal maturity in both oilfields.




Figure 9)  $C_{29}$  Sterane diagram in  $C_{32}$  hopane of Surmeh reservoir in Resalat and Salman oilfields.

The most important parameter of mature measurement by use of biomarkers is the rate of two epimerized shapes (20R, 20s) of Sterane  $\alpha\alpha$  which their rate is  $20s/(20s+20r)$ . The 20R figure is Sterane's  $\alpha\alpha\alpha$  bio form but when maturity is increasing turn to  $\alpha\alpha\alpha - 20s$ . When these two forms have their balanced, their ratio don't change, thus after this case, we can't register mature changes. In fact, mature changes have determined by the study of ratio change in  $20S/(20S+20R)$  in  $C_{29}$  Sterane (Justwanet al.,2006).

Regarding to the sample's location in chart maturity of Steranes in Resalat oilfield and

Surmeh layer shows good oil. Samples are located in upper part of oil window and near to window of the peak of thermal maturity and the situation implies on a suitable thermal maturity. Such condition illustrates a good thermal maturity in Surmeh layer in Resalat oilfield.

Also, according to the Figure 9 and this chart in Resalat and Salman oilfields in Surmeh layer thermal maturity lies between sub-mature to mature.




Figure 10) 20S/ (20S+20R) to  $\alpha\beta/(\alpha\beta+\alpha\alpha)$  of C29 Sterane isomers in samples from Surmeh source.

While maturity goes up Tm gradually would be disappeared and Ts increases. Unlike, Mortan to Hopane ratio, amount of Tm/Ts decrease. Tm/Ts amount is not a good factor in saline area because the primary Ts is high in samples from saline area (Justwanet al.,2006).

Thermal maturity in Salman and Arab layer is good and samples are in early stages of mature zone making use of Ts, Tm and C29 Sterane. X2 and X5 samples are located in the peak of this zone and have better condition while the others show a little poor quality.

The following describe thermal maturity in Resalat oilfield and Arab layer. Samples from this oilfield like Salman oilfield are located in

primary zone and are matured so thermal maturation is good. In comparison with the two other oilfields thermal maturation in Salman oilfield has better maturity and Arab layer has better maturity either. The reason why Salman oilfield is more mature is the depth which Surmeh layer is burdened in Salman oilfield which is deeper than Surmeh layer in Resalat oilfield.




Figure 11) chart of 20S/ (20S+20R) to  $\alpha\beta/(\alpha\beta+\alpha\alpha)$  of C29 Sterane isomer ratio, Surmeh source, Resalat and Salman oilfields.

Geological age of source rock has been assessed make use of “ $\delta^{13}C$ ” and C29 Sterane. The isotopes are in two forms, stable and unstable (radioactive). All isomer samples in this case imply on cretaceous and old and mature oil (Seifert and Moldowan, 1986, Table 4).

Table 1)  $\delta^{13}C$  parameter to sterane C29  $\beta\beta/(\beta\beta+\alpha\alpha)$ , Resalat and Salman oilfields.

Sample	$\delta^{13}C$ Sat	C29
Salman	-27.31	0.59
Salman	-26.98	0.62
Salman	-27.33	0.57
Salman	-27.19	0.51

Galimov chart could be used for aging the source rock (Galimov, 1973). In this chart, making use of samples analysis and the amount of  $\delta^{13}C$  for Asphaltine, Resins, whole oil and saturated oil, samples could be plotted on chart and characteristics of whole oil and source rock could be assessed (Fig. 12) .




Figure 12)  $\delta^{13}C$  to  $C_{29} / (\beta\beta + \alpha\alpha)$  sterane, Salman and Resalat oilfields.

Studies has been conducted by Galimov and his chart imply on the isotopic amount of carbon of hydrocarbon is related to age, the amount of isotopic carbon, the less aged the crust is the more weight the isotopes have. Regarding to table above and the line graph related to all samples have same source and they accumulate between -26 to -27 and the probable source rock maturity refers to beginning of oil window and its age is Cretaceous. Also, Y1 sample is better sample and shows better saturated carbon isotope (Table 5).

Table 5)  $\delta^{13}C$  Asph, Top oil, Resin, Sat, Aro fractions in Salman and Resalat oilfields.

Sample	$\delta^{13}C$ Asp %	$\delta^{13}C$ Res %	$\delta^{13}C$ Oil %	$\delta^{13}C$ Sat %	$\delta^{13}C$ Aro %
Resalat Y1	-26.0	-26.5	-26.7	-27.2	-26.5
Salman X5	-26.8	-26.3	-26.9	-27.2	-26.3

In this chart regarding to its saturated carbon isotope and top carbon isotope that are respectively -27 and -26.5 the X7 sample has the best condition. Resalat oilfield has the second best sample after Salman oilfield. Sample Y3 from Resalat oilfield and Surmeh layer has top carbon isotope -26.8 is better than X6 sample from Salman oilfield. As we know, the lighter the carbon isotope is the better condition it has and regarding to light hydrocarbon and the amount of them hydrocarbon from Salman and Resalat oilfields are heavy to moderate (Table 6).

Making use of changes in Sterane to Hopane ratio to  $C_{27}/C_{29}$  Sterane the geographic

locations of source rock could be defined. Also lithology of source rocks could be defined by this chart (Andrusevich et al., 2000).




Figure 13)  $C_{13}\delta$  chart for Asph, Top oil, Resin, Sat and Aro, Samples from Resalat and Salman oilfields.

Table 6)  $C_{13}\delta$  parameter for Asph, Top oil, Resin, Sat and Aro, Samples from Resalat and Salman oilfields, Surmeh reservoir.

Sample	Asphaltenes	Topped oil	Whole oil	Sat	Aro
Resalat Y3	-26.3	-26.8	-26.9	-26.8	-26
Salman X6	-26.1	-26.7	-26.8	-26.9	-26
Salman X7	-27	-26.7	-26.6	-27	-26

According to the Table 7 the chart for latitude could be illustrated and regarding to samples from upper latitudes have classic and in fact shale lithology while lower latitudes have non-clastic source rock and carbonate lithology.

Table 7)  $C_{27}/C_{29}$  steranes and Steranes/Hopanes ratios in Salman and Resalat oilfields samples.

Sample	$C_{27}/C_{29}$ Steranes	Steranes/hopanes
Salman X1	0.9	0.21
Salman X2	0.87	0.23
Salman X3	0.77	0.17
Salman X4	0.58	0.17
Resalat Y1	0.69	0.18
Resalat Y2	0.65	0.24

Firstly, the geographic location of Salman oilfield would be assessed. According to all charts and regarding to Sterane to Hopane volume is less than 0.3 and C27 Sterane to C29 is less than 1 it could be resulted that source rocks formed in lower latitude and are non-clastic rocks. Surmeh layer in Salman oilfield is like Arab layer and has carbonate originate (fig 14).

Secondly, Resalat oilfield would be investigated. According to studies Sterane to hopane ration is less than 0.24 and C27 to C29 Sterane below 0.7 is located in lower latitude (Fig.14).

Regarding to this, Salman and Resalat oilfields are near to each other and both have non-clastic or carbonate source rock.




Figure 14) Sterane to Hopane ratio against C27/C29 Sterane chart, Samples from Salman and Resalat oilfield.

#### 4- Conclusions

All geochemistry and biomarker parameters in Surmeh reservoir in Salman and Resalat oilfield like hydrocarbon quality, origin, organic matter and sedentary basin, organic and oil maturity grade in reservoir, hopane and sterane biomarkers, source rock age, latitude of source rock could be assessed making use of GC and GCMS and biomarkers and carbon isotopes. Regarding to all parameters, Surmeh reservoir in both Resalat and Salman oilfield produces same oil and this implies that there is lateral continuity in Surmeh reservoir.

In addition, the samples have same physical and chemical properties and carbon isotopic analysis

shows there is a unique source for both reserve. Finally, it is accepted that Resalat and Salman have same source rock there is no oil integration in the studied case.

#### References

- Andrusevich, V.E., Engel, M.H., Zumberge, J. E. 2000. Effects of paleo latitude on the stable carbon isotope composition of crude oils". *Geology*: 28, 847– 850.
- Bordenave, M. L. 1993. *Applied petroleum geochemistry*, Edition Technip, Paris, pp. 524.
- Connan, J., Cassau, A. M. 1980. Properties of gases and petroleum liquids derived from terrestrial kerogen at various maturation levels. *Geochimica et Cosmochimica Acta*: 44, 10–23.
- De Leeuw, J.W., Frewin, N.L., Van Bergan, P.F., Sinningh-Damste J.S., Collinson, M.E., 1995. Organic carbon as a paleo environmental indicator in marine realm. *Geological Society of London*: 83, 43–71.
- Didyk, B.M., Simoneit, B.R.T., Brasell, S.C., Eglinton, G. 1978. Organic geochemical indicators of paleo environmental conditions of sedimentation. *Nature*: 272, 216–222.
- Ekweozor, C.M., Okogun, J.I., Ekong, D.E.U. and Maxwell, J.R., 1979. Preliminary organic geochemical studies of samples from the Niger Delta (Nigeria) .1, Analyses of crude oils for Triterpenes. In: *Chemical Geology* vol. 27 P.11-28.
- Galimov, E. M. 1973. *Geochemistry of carbon and its applications in oil and gas exploration*, Moscow, Nedra, PP.383.
- Hunt, J.M. 1996. *Petroleum geochemistry and Geology*, 2<sup>nd</sup> edition. W.H, Freeman and company, New York. pp. 743.
- Justwan, H., Dahl, B., Isaksen, G.H. 2006. Geochemical characterization and genetic origin of oils and condensates in the South Viking Graben, Norway. *Marine and Petroleum Geology*: 23, 213–239.
- Kaufman, R.I, Ahmad, A.S., Elsinger, R.J. 1990. Gas chromatography as development and production tools for fingerprinting oils from individual reservoirs: application in the Gulf of Mexico, In: *GCSSEPM foundation ninth annual research conference proceedings*, pp.263–282.

- Moldowan, J.M., Dahl, J., Huizinga, B. J., Fago, F. J., Hickey, L. J., Peakman, T. M., Taylor, D.W. 1994. The molecular fossil record of oleanane and its relation to angiosperms. *Science*: 265, 768–771.
- Ourisson, G., Albrecht, P., Rohmer, M. 1984. The microbial origin of fossil fuels. *Scientific American*: 251, 44–51.
- Peters, K.E., Walters, C.C., Mold Owan, J.M. 2005. *The Biomarker Guide, Volume 2: Biomarkers and Isotopes in the Petroleum Exploration and Earth History*, Cambridge University Press; 2<sup>nd</sup> edition, PP. 684–700.
- Rabbani, A.R., 2013. *Petroleum geology and geochemistry of the Persian Gulf*. Tafresh: Tafresh University, Iran.
- Rabbani, A.R., Kotarbab, M. J., Baniadasa, A. R., Hosseinya, E., Wieclaw, D. 2014. Geochemical characteristics and genetic types of the crude oils from the Iranian sector of the Persian Gulf. *Organic Geochemistry*: 70, 29–43.
- Seifert, W. K., Moldowan, J.M. 1986. Use of biological markers in petroleum exploration, In: *Methods in Geochemistry and Geophysics*: 24, 261–90.
- Tissot, B.P., Welte, D.H. 1984. *Petroleum formation and occurrence*, 2<sup>nd</sup> Edition. Springer-Verlag, New York.
- Waples D. W., Machihara, T. 1991. Biomarker for geologist, a practical guide to the application of steranes and triterpenes in petroleum geology, AAPG Methods in exploration series 9. PP. 91.

**Received: 27 December 2015 / Accepted: 13 May 2016 / Published online: 15 May 2016**

#### **EDITOR-IN-CHIEF:**

##### ***Dr. Vahid Ahadnejad:***

Payame Noor University, Department of Geology, PO BOX 13395–3697, Tehran, Iran.  
E-Mail: edchief@jtethys.org

#### **EDITORIAL BOARD:**

##### ***Dr. Jessica Kind***

ETH Zürich Institut für Geophysik, NO H11.3, Sonneggstrasse 5, 8092 Zürich, Switzerland  
E-Mail: jessica.kind@erdw.ethz.ch

##### ***Prof. David Lentz***

University of New Brunswick, Department of Earth Sciences, Box 4400, 2 Bailey Drive Fredericton, NB E3B 5A3, Canada  
E-Mail: dlentz@unb.ca

##### ***Dr. Anita Parbhakar-Fox***

School of Earth Sciences, University of Tasmania, Private Bag 126, Hobart 7001, Australia  
E-Mail: anitap1@utas.edu.au

##### ***Prof. Roberto Barbieri***

Dipartimento di Scienze della Terra e Geoambientali, Università di Bologna, Via Zamboni 67 – 40126, Bologna, Italy  
E-Mail: roberto.barbieri@unibo.it

##### ***Dr. Anne-Sophie Bouvier***

Faculty of Geosciences and Environment, Institut des sciences de la Terre, Université de Lausanne, Office: 4145.4, CH-1015 Lausanne, Switzerland  
E-Mail: Anne-Sophie.Bouvier@unil.ch

##### ***Dr. Matthieu Angeli***

The Faculty of Mathematics and Natural Sciences, Department of Geosciences, University of Oslo  
Postboks 1047 Blindern, 0316 OSLO, Norway  
E-Mail: matthieu.angeli@geo.uio.no

##### ***Dr. Miloš Gregor***

Geological Institute of Dionys Stur, Mlynska Dolina, Podjavorinskej 597/15 Dubnicaná Vahom, 01841, Slovak Republic  
E-Mail: milos.gregor@hydrooffice.org

##### ***Dr. Alexander K. Stewart***

Department of Geology, St. Lawrence University, Canton, NY, USA  
E-mail: astewart@stlawu.edu

##### ***Dr. Cristina C. Bicalho***

Environmental Geochemistry, Universidade Federal Fluminense – UFF, Niterói–RJ, Brazil  
E-mail: ccbicalho@gmail.com

##### ***Dr. Lenka Findoráková***

Institute of Geotechnics, Slovak Academy of Sciences, Watsonova 45, 043 53 Košice, Slovak Republic  
E-Mail: findorakova@saske.sk

***Dr. Mohamed Omran M. Khalifa***

Geology Department, Faculty of Science, South  
Valley, Qena, 83523, Egypt  
E-Mail: mokhalifa@svu.edu.eg

***Prof. A. K. Sinha***

D.Sc. (Moscow), FGS (London). B 602,  
VigyanVihar, Sector 56, GURGAON 122011,  
NCR DELHI, Haryana, India  
E-Mail: anshuksinha@gmail.com